



# New Jersey

## Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,\* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
40th highest	44th highest	44th highest

\*This graph refers to females aged 15–19 years.




# Sexually transmitted diseases New Jersey

## Gonorrhea rates, 1982–1992



## Gonorrhea rates, 1992



## National ranking,\* 1992


Males – 29th highest    Females – 34th highest

\*National ranking refers to males and females aged 10–24 years.




## Chlamydia cases and rates,\* 1988–1992


Cases




Rates





Cases



Rates



## Acquired immunodeficiency syndrome (AIDS) cumulative cases, 1981–1992



\*Chlamydia cases and rates refer to males and females of all ages. Data were unavailable before 1991.




# Related risk behaviors,\* 1993

New Jersey

Had sexual intercourse (lifetime)




Had sexual intercourse with four or more people (lifetime)



Had sexual intercourse during the past three months (currently sexually active)



Used a condom during last sexual intercourse (among currently sexually active)



Used birth control pills during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



\*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 New Jersey Youth Risk Behavior Survey (risk behavior data).

